

The redshift periodicity of galaxies as a probe of the correctness of general relativity*

Valerio Faraoni

Department of Physics and Astronomy, University of Victoria

P.O. Box 3055, Victoria, B.C. Canada V8W 3P6

*Tel: [+]
604 721 8656, Fax: [+]
604 721 7715*

e-mail: faraoni@uvphys.phys.uvic.ca

Abstract

Recent theoretical work determines the correct coupling constant of a scalar field to the Ricci curvature of spacetime in general relativity. The periodicity in the redshift distribution of galaxies observed by Broadhurst *et al.*, if genuine, determines the coupling constant in the proposed scalar field models. As a result, these observations contain important information on the problem whether general relativity is the correct theory of gravity in the region of the universe at redshifts $z < 0.5$.

To appear in *General Relativity and Gravitation*

*This paper is based on a essay selected for an Honorable Mention by the Gravity Research Foundation, 1996.

The deep pencil-beam survey of the redshift distribution of galaxies at the Galactic poles by Broadhurst, Ellis, Koo and Szalay [1]–[3] has detected an apparent periodicity in the galaxy distribution with a period $128 h^{-1}$ Mpc (here $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$ is the present value of the Hubble parameter). The galaxy sample and the statistics used in [1] has been re-analyzed [4]–[6] and the result in [1] has been criticized. Additional data collected in the same directions confirmed the original results, supporting the idea that the peaks in the redshift distribution of galaxies are true features and not an artifact of inappropriate statistics [3]. Most of the peaks in the galaxy redshift distribution reported in [1] have been recognized and confirmed in [7]. The redshift survey at the Keck Telescope discussed in [8] also found evidence for strong clustering of galaxies in redshift space, although the redshift periodicity was not confirmed. If the periodicity in the redshifts of galaxies is real, one expects a similar phenomenon to occur also for the redshifts of Ly α absorption lines and for Mg II quasar absorption systems; evidence for periodicity in the redshift distribution of the Lyman- α forest was found in [9] but not in later observations [10, 11], and it was argued that it is not genuine [12]. The controversy is reminiscent of the old debate about the periodicity of the redshifts of quasars [13]–[16], or of similar claims for galaxies [17, 18]. Although many authors contend that the result in [1] is probably an artifact of incomplete sampling, it is generally recognized that further observations (expecially in directions different from the Galactic poles) are needed to definitely resolve the issue, and new surveys are planned or in progress [19, 8]. It is established that galaxies tend to cluster in sharp walls, leaving vast regions devoid of galaxies [20]–[22]; however, a single distinguished scale as found in [1] has not been confirmed. On the other hand, there is no doubt that, if confirmed, the claimed redshift periodicity would be of the outmost importance, and therefore it has drawn the attention of theoreticians [23]–[28]. In order to explain the periodicity found in [1]–[3], models were proposed in which clustering of galaxies in foam-like structures occurs at the predicted redshifts [29, 30]. A difficulty of these models is the implication that galaxies be approximately distributed on shells, of which we happen to be at the center, in conflict with the cosmological principle. A model in which the cosmological principle is explicitly violated was proposed recently [27], but it is non-viable. In fact, it assumes that the universe is represented by a closed Friedmann model dominated by a massless, minimally coupled, non self-interacting scalar field ψ . The energy density ρ and pressure P of this field are given by

$$\rho = P = \frac{1}{2} \nabla_\alpha \psi \nabla^\alpha \psi . \quad (1)$$

The stiff equation of state $P = \rho$ certainly does not describe our present, matter-dominated universe, in which the peculiar velocities of galaxies give a negligible con-

tribution to the pressure of the cosmological fluid ($P = 0$). A viable version of the model in [27] would necessarily have to include a mass and/or a non-minimal coupling for the scalar field. Such a model would be similar to the “oscillating universe” models introduced in [23, 24], in which the redshift periodicity of galaxies is an optical illusion rather than a real mass distribution. These other models are based on an oscillating scalar field non-minimally coupled to the Ricci curvature of spacetime. We show that if the redshift periodicity of the galaxy distribution is real, then the observations contain important information on the problem whether general relativity is the correct theory of gravity. This result arises from the observational determination of the coupling constant between the scalar field and the curvature [23, 24], and from recent theoretical work on the correct coupling in metric theories of gravity [31, 32].

In order to explain the observed distribution of galaxies, Morikawa [23] postulated a universe in which the dark matter is dominated by a massive, non-minimally coupled, scalar field. His model was shown to be non-viable by Hill, Steinhardt and Turner [24]. They proposed three alternative models involving oscillating physics: a) an oscillating “constant” of gravitation; b) oscillating atomic lines due to time variations in the fine structure constant, or in the electron mass; c) oscillating galactic luminosities. Model b) was ruled out by Sudarsky [25] on the basis of the Braginsky–Panov experiment. Model a) was shown to be compatible with biological records on the Earth [26], and has recently received renewed interest [28].

The common feature underlying the models under consideration is a spatially coherent, oscillating scalar field non-minimally coupled to the Ricci curvature R of spacetime. The Lagrangian density of the theory is given by¹

$$\mathcal{L} = \frac{1}{16\pi G_0} R - \frac{1}{2} \nabla_\mu \phi \nabla^\mu \phi - \frac{\xi}{2} R \phi^2 - V(\phi) + \mathcal{L}_m, \quad (2)$$

where G_0 is the present value of the gravitational “constant”, ξ is the coupling constant of the scalar field, $V(\phi)$ is its self-interaction potential, and $\mathcal{L}_m$ is the Lagrangian density of matter other than the scalar field. The non-minimal ($\xi \neq 0$) coupling seems to be a common feature of models unifying gravity with the other interactions and is an essential ingredient of many works on inflation. Two choices of ξ are most often found in the literature: $\xi = 0$ (“minimal coupling”) and $\xi = 1/6$ (“conformal coupling”). It must be emphasized that the correct value of the coupling constant is unknown, which

¹We use the metric signature $-+++$. The Riemann tensor is given in terms of the Christoffel symbols $\Gamma_{\mu\nu}^\rho$ by $R_{\mu\nu\rho}^\sigma = \Gamma_{\mu\rho,\nu}^\sigma - \Gamma_{\nu\rho,\mu}^\sigma + \dots$ and the Ricci tensor is $R_{\mu\rho} = R_{\mu\nu\rho}^\nu$. ∇_μ is the covariant derivative operator.

can only be regarded as a major piece of missing information regarding the physics of the scalar field. Recently, it was found [31] that if ϕ is a non-gravitational field in a curved spacetime, the Einstein equivalence principle (see [33] for a precise formulation and a discussion) forces the value of ξ to be $1/6$. This unintuitive result follows from the study of the Green’s function representation of the solutions of the Klein–Gordon equation

$$\square\phi - \xi R\phi - \frac{dV}{d\phi} = 0 , \quad (3)$$

and from the requirement that, in the neighbourhood of any point of spacetime, the propagation of scalar waves resembles more and more closely the propagation in flat spacetime as the size of the neighbourhood becomes smaller and smaller. This is the Einstein equivalence principle [33] applied to the propagation of scalar waves. The quoted result (later confirmed in [32]) holds for any metric and its derivation is completely independent of conformal transformations, or of the conformal structure of spacetime, or of the field equations of the theory of gravity [31]. If the nature of the field ϕ is gravitational (e.g. the scalar field of Brans–Dicke theory), the statement that its physics resembles *locally* the physics in flat spacetime goes beyond the Einstein equivalence principle, which is a statement regarding only *non-gravitational* physics [33] (see [34] for a discussion). In addition, the equation satisfied by a gravitational scalar field may be different from eq. (3).

Metric theories of gravity [33] (including general relativity) satisfy the Einstein equivalence principle. Therefore, if the correct theory of gravity describing the region of the universe at redshifts $z < 0.5$ observed by the survey in [1]–[3] is general relativity, or a metric theory in which the scalar field advocated by Hill, Steinhardt and Turner [24] is of non-gravitational origin, then the Einstein equivalence principle holds, and the coupling constant $\xi = 1/6$, as explained in [31].

Another prescription for ξ is found in a wide class of theories including Kaluza–Klein, higher derivative, supergravity and superstring-inspired theories. These theories are originally formulated in the “Jordan frame” (see [35] for definitions and the terminology) and are subsequently reformulated in the “Einstein frame” by means of a conformal transformation. The necessity and uniqueness of the conformal transformation were established for Brans–Dicke and Kaluza–Klein theories [36, 37] and later proved for a wider class of theories [35]. As a consequence of the conformal transformation to the Einstein frame, the conformally transformed scalar field is always minimally coupled to the Ricci curvature of spacetime² $\xi = 0$.

²To prevent a misunderstanding let us mention at this point that Einstein’s theory with a minimally

If the scalar field ϕ has a quantum nature, other prescriptions apply according to the nature of ϕ . If ϕ is a Goldstone boson, $\xi = 0$ [39]; if ϕ is the composite boson of the Nambu–Jona–Lasinio model, the value $\xi = 1/6$ was found in the large N approximation [40]; if $V(\phi) = \eta\phi^3/6$, then $\xi = 0$ [41], while $\xi \leq 0$ or $\xi \geq 1/6$ for Higgs scalar fields in the standard model [41]. The value of ξ in a quantum theory is also affected by renormalization [42]. In this paper, we limit ourselves to a *classical* scalar field ϕ . In fact, the universe at the epoch of galaxy formation was classical, and it is believed that also the scalar field driving inflation (with which the field ϕ has been identified [23]) can also be described classically, apart from the small quantum perturbations generating seeds for the formation of structures.

It is significant that all the viable models proposed in order to describe the observed distribution of galaxies are based on a classical, oscillating, scalar field non-minimally coupled to gravity, and that the value of the constant ξ is a parameter of the models, which can be determined by the observations. Typically, the period of the oscillations in redshift of the galaxy distribution ($128 h^{-1}$ Mpc) determines the mass of the scalar field, while the amplitude of the oscillations determines the coupling constant ξ [23, 24]. It is also possible to employ a massless ϕ ; a variation of the model with oscillating gravitational “constant” predicts $\xi \approx 6.4 \cdot 10^3$ [24]. An unambiguous fitting of the observational data with a value of ξ significantly different from $1/6 \simeq 0.166$ implies that general relativity is not the correct theory of gravity in the observed region of the universe $z < 0.5$, because the Einstein equivalence principle is violated. This would be a remarkable result, since Solar System experiments seem to indicate that general relativity is the most plausible theory of gravity [33]. There remains two possibilities:

- i) ϕ is a gravitational field in a metric theory of gravity other than general relativity: in this case the Einstein equivalence principle and the prescription $\xi = 1/6$ do not apply to ϕ ;
- ii) gravity is described by a non-metric theory of gravitation, in which the Einstein equivalence principle (and consequently the prescription $\xi = 1/6$) does not necessarily hold.

The latter possibility is regarded as highly unlikely, since there are good reasons to believe that the correct theory of gravity is a metric theory [33]. If case i) is appropriate, the observational determination of the value of ξ is not sufficient to single out the correct theory of gravity, but can nevertheless provide a important hint. If one of the models in [24] is correct, then $\xi \neq 0$ and the correct (low-energy) theory of gravity is *not*

coupled scalar field and Einstein’s theory with a conformally coupled scalar field are also conformally equivalent to each other (cf. e.g. [38]).

one formulated in the Einstein frame: this would exclude Brans–Dicke, Kaluza–Klein, supergravity and superstring theories. In addition, a reliable determination of ξ , which is otherwise unknown, is regarded as a remarkable progress in the physics of the scalar field.

The spatial coherence of the scalar field ϕ has been related to inflation [24], and the scalar field ϕ has been explicitly identified with the inflaton [23]. If ϕ is to be identified with the field driving inflation, the value of the coupling constant has a profound effect on the success of the inflationary scenario under consideration [34]. Since the proposed inflationary scenarios are currently under test using observations of the cosmic microwave background, a reliable measurement of the coupling constant ξ would give further hints on the correctness of some of the proposed scenarios for inflation. However, it is to be stressed that the region of the universe under study (redshifts $z < 0.5$) does not correspond to the very early universe, which is observationally inaccessible, but it is well within the range of present astronomy.

Ultimately, the problems whether one of the oscillating universe models is correct and whether it is possible to measure the coupling constant ξ are meaningful only if the redshift periodicity discovered in [1]–[3] is genuine. Since general relativity has been successfully tested in the Solar System [33], the belief that it is valid at least in the region of the universe at redshifts $z < 0.5$ has a firm justification, and one could go as far as considering this fact, in conjunction with the suggested values of ξ ($\xi = 10$ [23], $\xi = 6.4 \cdot 10^3$ [24], $\xi = 6.267$ or other values [28]), as an argument against the authenticity of the redshift periodicity reported in [1]–[3]. While we do not support this extreme point of view, we believe that the possibility of measuring the coupling constant ξ and its implications for the determination of the correct theory of gravity provide new and compelling motivation for resolving the issue of the redshift periodicity of galaxies. To understand the large-scale clustering of galaxies, new pencil beams have been analyzed and new surveys are being carried on [19, 22].

Acknowledgments

The author acknowledges F.I. Cooperstock, C.J. Pritchett and R. De Propris for stimulating discussions, and a referee for suggestions leading to improvements in the manuscript. This work was supported by the NATO Advanced Fellowships Programme through the National Research Council of Italy (CNR).

References

- [1] T.J. Broadhurst, R.S. Ellis, D.C. Koo and A.S. Szalay, *Nature* 343 (1990) 726.
- [2] D.C. Koo *et al.*, in ASP Conf. Ser. 51, *Observational Cosmology*, ed. by G. Chincarini *et al.* (ASP, S. Francisco), p. 112.
- [3] A.S. Szalay *et al.*, *Proc. Nat. Acad. Sci.* 90 (1993) 4853.
- [4] N. Kaiser and J.A. Peacock, *Astron. J.* 379 (1991) 482.
- [5] C. Park and J.R. Gott, *Mon. Not. R. Astr. Soc.* 249 (1991) 288.
- [6] A. Dekel, G.R. Blumenthal, J.R. Primack and D. Stanhill, *Mon. Not. R. Astr. Soc.* 257 (1992) 715.
- [7] C.N.A. Willmer *et al.*, *Astrophys. J.* 437 (1994) 560.
- [8] J.C. Cohen, D.W. Hogg, M.A. Pahre and R. Blandford, *Astrophys. J. Lett.* 462 (1996) L9.
- [9] Y. Chu and X. Zhu, *Astron. Astrophys.* 222 (1989) 1.
- [10] J.G. Bartlett, R. Esmailzadeh and L.J. Hall, UC Berkeley preprint (1990).
- [11] D. Tytler, J. Sandoval and X.M. Fan, *Astrophys. J.* 405 (1993) 57.
- [12] D. Scott, *Astron. Astrophys.* 242 (1991) 1.
- [13] K.G. Karlsson, *Astr. Astrophys.* 13 (1971) 333.
- [14] S. Depaquit, J.C. Pecker and J.P. Vigier, *Astr. Nachr.* 306 (1985) 7.
- [15] H. Arp, H.G. Bi, Y. Chu and X. Zhu, *Astr. Astrophys.* 239 (1990) 33.
- [16] A. Holba *et al.*, *Astrophys. Sp. Sci.* 222 (1994) 65.
- [17] W.G. Tifft, *Astrophys. J.* 206 (1974) 38.
- [18] B.N.G. Guthrie and W.M. Napier, *Mon. Not. R. Astr. Soc.* 253 (1991) 533.
- [19] D. Fabricant, E. Hertz and A. Szentgyorgy, *Proc. SPIE*, 2198 (1994); M.J. Geller, J. Roy. Astr. Soc. Can. 88 (1994) 283; G. Vettolani *et al.*, in *Proc. 9th IAP Meeting, Cosmic Velocity Fields*, ed. by F.R. Bouchet and M. Lachièze-Rey (Editions Frontieres, Gif sur Yvette), p. 523.

- [20] M.J. Geller and J.P. Huchra, *Science* 246 (1989) 897.
- [21] L.N. da Costa *et al.*, *Astrophys. J.* 424 (1994) L1.
- [22] C. Bellanger and V. de Lapparent, *Astrophys. J.* 455 (1995) L103.
- [23] M. Morikawa, *Astrophys. J. Lett.* 362 (1990) L37; *Astrophys. J.* 369 (1991) 20.
- [24] C.T. Hill, P.J. Steinhardt and M.S. Turner, *Phys. Lett. B* 252 (1990) 343.
- [25] D. Sudarsky, *Phys. Lett. B* 281 (1992) 281.
- [26] P.D. Sisterna and H. Vucetich, *Phys. Rev. Lett.* 72 (1994) 454.
- [27] P. Budinich, P. Nurowski, R. Raczka and M. Ramella, *Astrophys. J.* 451 (1995) 10; P. Budinich and R. Raczka, *Found. Phys.* 23 (1993) 225; *Europhys. Lett.* 23 (1993) 295.
- [28] M. Salgado, D. Sudarsky and H. Quevedo, *Phys. Rev. D* 53 (1996) 6771.
- [29] R. van de Weygaert, *Mon. Not. R. Astr. Soc.* 249 (1991) 159.
- [30] S. Ikeuchi and E.L. Turner, *Mon. Not. R. Astr. Soc.* 250 (1991) 519.
- [31] S. Sonogo and V. Faraoni, *Class. Quant. Grav.* 10 (1993) 1185.
- [32] A.A. Grib and E.A. Poberii, *Helv. Phys. Acta* 68 (1995) 380.
- [33] C. M. Will, *Theory and Experiment in Gravitational Physics*, revised edition (Cambridge Univ. Press, Cambridge, 1993).
- [34] V. Faraoni, *Phys. Rev. D* 53 (1996) 6813.
- [35] G. Magnano and L.M. Sokolowski, *Phys. Rev. D* 50 (1994) 5039.
- [36] Y.M. Cho, *Phys. Rev. Lett.* 68 (1992) 3133.
- [37] L. Bombelli *et al.*, *Nucl. Phys. B* 289 (1987) 735; L.M. Sokolowski and Z. Golda, *Phys. Lett. B* 195 (1987) 349; L.M. Sokolowski, *Class. Quant. Grav.* 6 (1989) 59; 2045.
- [38] S. Deser, *Phys. Lett. B* 134 (1984) 419.
- [39] M.B. Voloshin and A.D. Dolgov, *Sov. J. Nucl. Phys.* 35 (1982) 120.
- [40] C.T. Hill and D.S. Salopek, *Ann. Phys. (NY)* 213 (1992) 21.
- [41] Y. Hosotani, *Phys. Rev. D* 32 (1985) 1949.
- [42] M. Reuter, *Phys. Rev. D* 49 (1994) 6379; L. Parker and D.J. Toms, *Phys. Rev. D* 32 (1985) 1409.